

Dynamics of Collapse of a Confined Bose Gas

L.P. Pitaevskii

Department of Physics, Technion, 32000 Haifa, Israel

and

Kapitza Institute for Physical Problems, 117454 Moscow, Russia

(April 11, 1996)

Abstract

Rigorous results on the nonlinear dynamics of a dilute Bose gas with a negative scattering length in an harmonic magnetic trap are presented and sufficient conditions for the collapse of the system are formulated. By using the virial theorem for the Gross-Pitaevskii equation in an external field we analyze the temporal evolution of the mean square radius $I = \int r^2 |\psi|^2 dV$ of the gas cloud. In the 2D case this quantity undergoes harmonic oscillations with frequency $2\omega_0$. It implies that for a negative value of the energy of the system, the gas cloud will collapse after a finite time interval. For positive energy the cloud collapses if the initial conditions correspond to a large enough amplitude of oscillations. Stable oscillations with a small amplitude are possible. In the 3D case the system also collapses after a finite time for a state with negative energy. A stringent condition for the collapse is also derived.

The recent observation of Bose-Einstein condensation of atoms of ^{87}Rb [1], ^7Li [2] and ^{23}Na [3] confined in magnetic traps has opened a new important field of investigation of quantum phenomena. In particular, a new feature of these systems is that they are significantly inhomogeneous so one can directly observe effects of quantum uncertainty on a macroscopical scale. The formal evidence of such a situation is that dynamics of these non-uniform gases at low temperatures can be properly described by a non-linear equation for the condensate wave function $\psi(\mathbf{r}, t)$ [4], [5]:

$$i\hbar \frac{\partial}{\partial t} \psi(\mathbf{r}, t) = \left(-\frac{\hbar^2 \nabla^2}{2m} + U(\mathbf{r}) + g |\psi(\mathbf{r}, t)|^2 \right) \psi(\mathbf{r}, t). \quad (1)$$

Here U is the confining potential and $g = \frac{4\pi\hbar^2 a}{m}$, where a is the s-wave scattering length. This equation is quite classical in its merit but contains explicitly the Planck's constant $\hbar$. (We will put $\hbar = m = 1$ below.)

Experimental evidences of BEC in magnetically trapped alkali vapour stimulated recent numerical investigations of Eq. (1) in static, linear and nonlinear regimes. [6] - [10]

The physical situation is very different for ^{87}Rb , ^{23}Na on one side and ^7Li on the other. In the first case $a > 0$ and the homogeneous state of the gas is stable. For ^7Li the scattering length a is negative ($a = -1.43 \times 10^{-7}$ cm), which corresponds to an effective attraction, and the gas possesses negative compressibility. The uniform state in this case is unstable and the gas can undergo a "collapse" which is an unrestricted contraction. (The idea of such a collapse was first introduced by V.Zacharov for a formally similar system of Langmuir waves [11].) Fetter however has recently shown [7], using a variational approach, first employed by Baym and Pethick [6], that a system of a finite number N of atoms in a trap can be in a metastable state for N below some critical number N_c . The same conclusion has been obtained by Ruprecht et al. [8]. Surprisingly the number of atoms reached in the experiment [2] appears significantly larger than the theoretical estimate of N_c .

In such an uncertain situation it is important to understand better the conditions and dynamics of the collapse.

In this paper we will present some rigorous results relative to the behaviour of the solution

of Eq. (1). Particularly we will formulate a sufficient condition for collapse in the gas. We will analyze the temporal evolution of the quantity $I = \int r^2 |\psi|^2 dV$, which is proportional to the average square of the radius of the gas cloud. This "virial theorem" has been used by Vlasov et al. to analyze the problem of the collapse, described by Eq.(1), in the absence of the confined field [12]. (See also more recent papers [13] - [17] and references therein.)

Note first of all that Eq. (1) provides conservation laws for the energy E of the system:

$$E = \int \left(\frac{1}{2} |\nabla \psi|^2 + \frac{g}{2} |\psi|^4 + U |\psi|^2 \right) dV \equiv E_0 + \int U |\psi|^2 dV \quad (2)$$

and for the number of atoms $N = \int |\psi|^2 dV$. The last conservation law is expressed by the continuity equation:

$$\frac{\partial |\psi|^2}{\partial t} + \frac{\partial j_i}{\partial x_i} = 0. \quad (3)$$

The equation for the momentum density in the presence of an external field can be written in the form:

$$\frac{\partial j_i}{\partial t} + \frac{\partial \Pi_{ik}}{\partial x_k} = - |\psi|^2 \frac{\partial U}{\partial x_i}. \quad (4)$$

We will not write explicitly the well known expressions for the currents j_i and Π_{ik} and we note only that in 2D and 3D we have an exact identities

$$\int \Pi_{ii} dV = 2E_0 \quad (5)$$

and

$$\int \Pi_{ii} dV = 3E_0 - \frac{1}{2} \int |\nabla \psi|^2 dV \quad (6)$$

respectively.

Repeating differentiation of I with respect to t and integrating by parts one gets:

$$\frac{dI}{dt} = 2 \int x_i j_i dV, \quad \frac{d^2 I}{dt^2} = 2 \int \Pi_{ii} dV - 2 \int x_i \frac{\partial U}{\partial x_i} |\psi|^2 dV. \quad (7)$$

We first analyze the two-dimensional motion of the gas in an isotropic cylindrical trap with a confining potential $U = \frac{\omega_0^2 r^2}{2}$, $r^2 = x^2 + y^2$. In this case Eq. (7) takes a surprisingly simple form:

$$\frac{d^2 I}{dt^2} = -4\omega_0^2 I + 4E. \quad (8)$$

whose general solution is of the obvious form:

$$I = A \cos(2\omega_0 t + \gamma) + E/\omega_0^2. \quad (9)$$

It is worth noting that the parameters A , α and E of the solution (9) can be calculated using the initial distribution $\psi(\mathbf{r}, 0)$ with the help of Eq.(2) and the first equality in Eq.(7).

Eq. (8) describes harmonic pulsations of I with constant frequency $2\omega_0$ in spite of the nonlinear character of Eq.(1). (It is possible to show that small axial symmetric oscillations of the system have the frequency $2\omega_0$ [18].) Furthermore, one can see that for a negative value of the energy the positive quantity I approaches zero after a finite time smaller than $\pi/2\omega_0$. This means complete contraction of the cloud. On the contrary, for a positive value of the energy pulsations with a small enough amplitude $A < E/\omega_0^2$ do not result in a collapse [19]. (See also the discussion in the concluding part of the paper.)

Thus increasing the energy improves the stability of the cloud. For example an increase in the stability has been noted in [20] in the presence of a vortex line.

Equation (8) has the same form both for attractive and repulsive interactions. In the latter case however the collapse does not occur. The confining potential energy $\omega_0^2 I/2$ is then always smaller than E . This inequality ensures that the collapse cannot be reached.

One can suggest that under the experimental conditions some motion of the superfluid gas exists that increases the energy of the system. This means that the cloud is not in an equilibrium but rather in a nonstationary stable state.

Results for the 3D case are not so conclusive. Consider an anisotropic harmonical trap with a confining potential of the form $U = \frac{\alpha_{ik} x_i x_k}{2}$. One gets from (7) and (6):

$$\frac{d^2 I}{dt^2} = 6E - 5 \int \alpha_{ik} x_i x_k dV - 2K, \quad (10)$$

where $K = \frac{1}{2} \int |\nabla \psi|^2 dV$ is the kinetic energy. Eq. (10) gives the inequality:

$$\frac{d^2 I}{dt^2} + 5\omega_m^2 I - 6E < 0, \quad (11)$$

where ω_m^2 is the smallest eigenvalue of the α_{ik} matrix. We can prove again that the system collapses after a finite time for a negative E . Consider the auxiliary equation

$$\frac{d^2 I}{dt^2} + 5\omega_m^2 I - 6E = 0. \quad (12)$$

For negative E the quantity I reaches, according to this equation, the value zero after a time smaller than $\frac{\pi}{\sqrt{5}\omega_m}$. The negative term in the left hand side of Eq.(10) accelerates this process for the same initial conditions. The same situation takes place for a positive energy for a large enough amplitude.

One can get more rigorous sufficient conditions of the collapse using the powerful method developed in [13]- [16]. In this method an important part is played by two inequalities imposed on the kinetic energy K [13], [15]. The first one is the uncertainty relation between I and K :

$$IK \geq \frac{9}{8}N^2. \quad (13)$$

The second inequality can be written as

$$\frac{1}{2} \int |\psi|^4 dV \leq \beta N^{1/2} K^{3/2}, \beta = 0.0575. \quad (14)$$

Combining (13) and (14) one immediately gets an inequality for the energy E as a function of K :

$$E(K) > K + \frac{9}{16}\omega_m^2 N^2 K^{-1} - \beta |g| K^{3/2} N^{1/2}. \quad (15)$$

This equation demonstrates a remarkable resemblance to the expression for $E(d^{-2})$ which has been obtained by Fetter using a wave function of the form $\psi = C \exp(-r^2/2d^2)$. (See Eq.(5) in [7].) For such a function the kinetic energy is proportional to d^{-2} . It easily checked that $E(d^{-2})$ (expressed as a function of K) is larger than the right hand side of (15). The equation (15) permits us to formulate the stringent collapse conditions we are looking for. Introducing dimensionless variables ϵ and κ defined by

$$E = \frac{3}{4}N\omega_m\epsilon, K = \frac{3}{4}N\omega_m\kappa, \quad (16)$$

we can rewrite this equation as:

$$\epsilon > \kappa + \kappa^{-1} - 1.18 \frac{2}{3} \sigma \equiv f(\kappa, \sigma), \quad (17)$$

where σ is a dimensionless parameter

$$\sigma = \sqrt{2/\pi} N |a| \omega_m^{1/2} \quad (18)$$

The parameter σ is analogous to the parameter σ defined in [7].

Behaviour of the function $f(\kappa)$ depends on the value of σ . For $\sigma \geq \sigma_c = 0.454$ the curve is a monotonous one (the curve (a) in Fig.1). Let the point ϵ^* , κ^* to be an intersection of the curve $\epsilon = f(\kappa)$ and the straight line $\epsilon = \kappa/3$. Let us to be

$$E < E^* = \frac{3}{4} N \omega_m \epsilon^* > 0. \quad (19)$$

Then the kinetic energy $K > K^* = 3E^*$ for any t . This means that equation (10) leads to the inequality

$$\frac{d^2 I}{dt^2} + 5\omega_m^2 I - 6(E - E^*) < 0, \quad (20)$$

This formula obviously demonstrates that at condition (19) we get the collapse.

We shall now discuss the case where $\sigma < \sigma_c$. The curve $\epsilon = f(\kappa)$ has in this case a minimum ϵ_m and a maximum ϵ_M (the curve (b) in Fig.1). Let us consider an initial state, where

$$\epsilon_m < \epsilon < \epsilon_M, \kappa_1 < \kappa < \kappa_2. \quad (21)$$

Since any solution of (1) evolves continuously, the kinetic energy κ will stay in the same range at every time. The uncertainty relation (13) gives then

$$I > \frac{3N}{2\omega_m \kappa_2}. \quad (22)$$

Thus the mean square radius stays restricted and collapse of the cloud as a whole is impossible.

If $E < E^*$ the cloud will collapse as in the previous case. The evolution of the cloud for $E > E^*$ and K out the interval (21) demands additional investigation.

The method we explored is also useful in the two-dimensional case. In this case instead of (13) and (14) we have

$$IK \geq \frac{1}{2}N^2, \frac{1}{2} \int |\psi|^4 dV \leq \beta NK, \beta = 0.17. \quad (23)$$

We introduce dimensionless variables ϵ and κ defined by:

$$E = \frac{1}{2}N\omega_m\epsilon, K = \frac{1}{2}N\omega_m\kappa. \quad (24)$$

Then we obtain the following inequality for the energy:

$$\epsilon > \kappa(1 - N/N_c) + \kappa^{-1} \equiv g(\kappa, N), \quad (25)$$

where $N_c = 1/\beta$ | g | The shape of the right hand side of this inequality is different for $N > N_c$ and for $N < N_c$. (See curves (a) and (b) in Fig.2 respectively.) In the latter case function g tends to $+\infty$ as κ tends both to 0 or ∞ . Hence the kinetic energy for a given value E stays restricted by some values κ_1 and κ_2 and the total collapse is impossible. This means that for $N < N_c$ an arbitrary initial function $\psi(\mathbf{r}, 0)$ automatically ensures such initial conditions for Eq. (8) that the collapse to be absent. In the opposite case, when $N > N_c$, it is impossible to draw by this method any general conclusions.

I would like to thank S. Stringari for various enlightening discussions and S. Giorgini for useful remarks. I am grateful for hospitality to the Department of Physics of the University of Trento where this paper has been completed.

REFERENCES

- [1] M.N. Anderson, J.R. Ensher, M.R. Matthews, C.E. Wieman, and E.A. Cornell, *Science* **269**, 198 (1965).
- [2] C.C. Bradley, C.A. Sacket, J.J. Tollet, and R.G. Hulet, *Phys. Rev. Lett.* **75**, 1687 (1995).
- [3] K.B. Davis, M.-O. Mewes, M.R. Andrews, N.J. van Druten, D.S. Durfee, D.M. Kurn, and W. Ketterly, *Phys. Rev. Lett.* **75**, 3969 (1995).
- [4] E.P. Gross, *Nuovo Cimento* **20**, 454 (1961); *J. Math. Phys.* **46** 137 (1963).
- [5] L.P. Pitaevskii, *Sov. Phys. JETP* **13**, 451 (1961).
- [6] G. Baym and C. Pethick, *Phys. Rev. Lett.* **76**, 6 (1996).
- [7] A. Fetter, (unpublished).
- [8] P.A. Ruprecht, M.J. Holland, K. Burnett, and M. Edwards, *Phys. Rev. A* **51**, 4704 (1995).
- [9] P.A. Ruprecht, M. Edwards, and K. Burnett, (unpublished).
- [10] M. Holland and J. Cooper, *Phys. Rev. A* **53**, R1954 (1996).
- [11] V.E. Zakharov, *Sov. Phys. JETP* **35**, 908 (1972).
- [12] S.N. Vlasov, V.A. Petrishcev, and V.I. Talanov, *Izv. Vyssh. Uchebn. Zaved., Radiofiz.* **12**, 1353 (1970).
- [13] M.I. Weinstein, *Commun. Math. Phys.* **87**, 567 (1983).
- [14] E.A. Kuznetsov, A.M. Rubenchik, and V.E. Zakharov, *Phys. Rep.*, **142**, 103 (1986).
- [15] S.K. Turitsyn, *Phys. Rev. E* **47**, R13 (1993).
- [16] E.A. Kuznetsov, J.J. Rasmussen, K. Rypdal, and S.K. Turitsun, *Physica D* **87**, 273 (1995).

- [17] P.M. Lushnikov, JETP Lett., **62**, 461 (1995).
- [18] S. Stringari, (private communication).
- [19] One must be careful about interpretation of these oscillations. The present considerations cannot exclude the possibility of a "partial collapse" of the denser part of the cloud. Collapsed atoms will escape the trap and the system will fit itself to a more stable state. The occurrence of a singular point where the density formally tends to infinity is also possible.
- [20] F. Dalfovo and S. Stringari, Phys. Rev. A. **53**, (1996).

FIGURES

FIG. 1. The function $f(\kappa)$ defined in (17) for $\sigma = 0.51 > \sigma_c$ (curve (a)) and for $\sigma = 0.29 < \sigma_c$ (curve (b)). Admitted states of the system are above the bold curve. The dashed line is $\epsilon = \kappa/3$.

FIG. 2. The function $g(\kappa)$ defined in (25) for $N = 0.4N_c$ (curve (a)) and for $N = 1.4N_c$ (curve (b)). Admitted states of the system are above the corresponding curve.